

ATOLL RESEARCH BULLETIN

No. 269

THE INVERTEBRATES OF GALETA REEF (CARIBBEAN PANAMA):
A SPECIES LIST AND BIBLIOGRAPHY

BY

JOHN CUBIT AND SUELYNN WILLIAMS

ISSUED BY

THE SMITHSONIAN INSTITUTION

WASHINGTON, D. C., U.S.A.

SEPTEMBER 1983

CONTENTS

	page
Abstract	1
Introduction	1
Explanation of Listings	5
Porifera	7
Coelenterata	8
Platyhelminthes	11
Sipuncula	12
Annelida	12
Arthropoda	19
Mollusca	28
Ectoprocta (Bryozoa)	38
Brachiopoda	38
Echinodermata	38
Chordata	40
Acknowledgments	41
Literature Cited	42

THE INVERTEBRATES OF GALETA REEF (CARIBBEAN PANAMA)
A SPECIES LIST AND BIBLIOGRAPHY

by John Cubit¹ and Sue Lynn Williams²

ABSTRACT

The fringing coral reef at Galeta Point, on the Caribbean Coast of Panama, has been under intensive study for approximately ten years. The combined studies of the reef represent approximately 5,000 person hours of field sampling and have documented the presence of approximately 775 species of invertebrates. A list of these species is presented here together with annotations to all known sources of information pertaining to the occurrence of these species on Galeta Reef. For most species these records are the southernmost in the Caribbean Sea.

INTRODUCTION

The following is an inventory of the invertebrates that have been found on Galeta Reef, Panama, together with all known sources of information pertaining to the occurrence of these species on the reef. At present such data are lacking for Caribbean reefs, but are needed to formulate rational policies of resource management and to serve as baseline information for measuring natural and man-caused changes of the environment (FAO, 1969). This need was recently reiterated by the United Nations Environment Program (UNEP, 1981). We have compiled this list from published papers, unpublished reports, personal communications, our own observations, and the set of specimens in the reference collection of the Galeta Point Marine Laboratory. This listing has been restricted only to those invertebrates studied on, or

-
1. Smithsonian Tropical Research Institute, Box 2072, Balboa, Panama.
 2. Present address: 9 Second Street, Route 4, West Riverside, Missoula, Montana 5980 .

Manuscript received May 1982 -- Eds.

collected from, Galeta Reef itself; it does not include species reported from nearby areas or the general Caribbean coast of Panama.

Galeta Reef has been under intensive investigation for approximately ten years. Most of the reef studies have been monitoring surveys designed to determine the species composition of the reef and to document temporal changes in the distribution and abundance of the biota of the reef.

This reef was established as a biological reserve in the late 1960s. Intensive monitoring of the reef biota was started by Charles Birkeland and others in 1970, beginning with a three-year program




Figure 1. Map of Caribbean Sea showing the location of Galeta Point (9°24'18"N, 79°51'48.5"W) (marked by an arrow at lower left).




Figure 2. Schematic cross section of the reef at Galeta Point. The reef platform (10 to 300 m wide) is in shallow water (less than 40 cm deep), and is occasionally exposed above the level of the tide (shown as a dashed line). The seaward edge of the reef has various profiles (shown in the figure as the solid and dotted lines) which terminate at 3 to 10 m depth at a base of sand or coral rubble. The reef platform encloses a lagoon and channels and is backed on the landward side by mangroves and sand beaches.

supported by the U.S. Environmental Protection Agency (see Birkeland et al., 1976). Subsequently the monitoring was expanded to include physical factors in a set of projects supported by the Smithsonian Institution Environmental Sciences Program. In addition, the reef has been studied by a number of visiting investigators. Overall, the information in this inventory has been derived from the efforts of approximately 50 people who have conducted a variety of studies, of variable duration, during the period of 1969 to 1980. In all, we estimate that this represents approximately 5,000 person-hours of sampling effort of reef invertebrates. The presence of approximately 775 species of macroinvertebrates has been documented in these investigations of the reef. In these studies certain taxonomic and ecological groups have received more attention than others. For instance, the gastropods, polychetes, and certain groups of decapods have been more thoroughly collected, while the sponges, ascidians, bryozoans, and smaller crustaceans have not. There is also more information regarding those invertebrates that are major occupiers of primary substratum on the shallow reef platform than for those species that are mobile or occur on the subtidal reef slope and sand bottom.

Description of the Reef

The Galeta Reef is in the southwestern Caribbean Sea ($9^{\circ}24'18''N$, $79^{\circ}51'48.5''W$). The Galeta Point Marine Laboratory, situated directly on the reef platform, is the southernmost marine laboratory in the Caribbean (see map, Fig. 1). The reef is near the western end of a system of fringing reefs that stretch for approximately 250 km along the Caribbean coast of Panama. The majority of these reefs are now at their post-climax stage, having passed through their period of most active coral growth more than 2,000 years ago. The general structure of this type of reef is of a broad platform, bordered on the landward side by mangroves, and often enclosing a lagoon. On the seaward side, the platform slopes into 3 to 10 m of water before reaching a sand bottom (Macintyre and Glynn, 1976; Fig. 2). The surface of the reef platform is at the lower level of the intertidal zone. The range of the tides on this coast is only about 30 cm, so the reef platform is never deeply under water. Wave action tends to maintain water over the reef even during the lowest low tides; however, during calm weather the reef platform may be exposed above water level for long periods of the day, subjecting the biota of the platform to extremes of desiccation, solar radiation, high temperatures, rainfall, and predation by shorebirds. There are approximately 30-40 such instances of reef exposure each year, varying in duration from 1 to 14 hours.

Other than the extreme low tides, the reef is subject to relatively little disturbance. Galeta Point is south of the Caribbean "hurricane belt" (the area traversed by hurricanes and tropical storms) (Neumann et al., 1978), and is thus protected from this source of periodic disruption to which most other Caribbean reefs are exposed. Salinity and seawater temperature have been monitored on this reef for approximately seven years: there is no evidence that the subtidal biota are ever exposed to extremes of any of these factors. Mean seawater temperature is approximately $29^{\circ}C$ with the range of variation confined to plus or minus a few degrees (except during the extreme low tides previously mentioned, when temperatures in shallow pools of standing water on the reef platform may reach $35^{\circ}C$ to $40^{\circ}C$). Salinities average approximately 33 ppt, also with little variation (except for those organisms above water level which may be exposed to heavy rains).

Most of the hard substrata of the reef platform are covered with algae, primarily the two red algae Acanthophora spicifera (Vahl) Borgesen and Laurencia papillosa (Forsskal) Greville. The seagrass Thalassia testudinum Koenig and Sims occupies much of the area of loose sediment. The biota of the subtidal reef slope consists of various species of foliose and crustose coralline algae, some fleshy algae, and live corals. Most of the species in this inventory were collected from the reef platform.

Explanation of Listings

All entries on this list are followed by notations which refer to the following: (1) published information regarding the occurrence of the species on Galeta Reef, (2) to whom to attribute the determination of the species, and (3) the presence of voucher specimens in the Galeta laboratory reference collection. Because of the large number of entries, we have had to abbreviate these notations. The meanings of the abbreviations are listed at the end of the inventory and follow this form:

1. Multiple capital letters are the initials of those persons who have made the species determinations of the specimens collected from Galeta Reef.
2. The single capitals B, D, F, and G refer to appendices in Birkeland et al. (1976), the single most comprehensive survey of Galeta Reef.
3. E73-E76 refer to reports of the Environmental Sciences Program for the years numbered.
4. Lower case letters denote references listed under Literature Cited.
5. + signifies voucher specimens of the species are present in the reference collection at the Galeta Point Marine Laboratory.

To aid users of this list and to avoid compounding errors, we have noted any variations in nomenclature and spelling among the sources. We refer the user to the sources in the annotations for the authors of the species.

Abbreviations Used

SPECIES DETERMINATIONS

AR	Amada A. Reimer
BM	B. Macurda
CAC	C. A. Child
CB	Charles Birkeland
CC	C. Cutress
DLM	David L. Meyer
DLW	Diana L. Werder
DS	Diane Stoecker
EK	Eugene Kaplan
GLH	Gordon L. Handler
HD	Hugh Dingle

IT	Ina Tumlin
JC	John Cubit
JD	Judith Dudley
JH	Janet Haig
JR	Joseph Rosewater
JRY	Joyce Redemske Young
JS	James Stames
JSG	John S. Garth
JW	J. Wells
JWP	J. W. Porter
KBM	Kaniaulono B. Meyer
KS	Ken Sebens
KR	Klaus Ruetzler
LGA	Lawrence G. Abele
PAA	Peter A. Abrams
PG	Peter W. Glynn
RB	R. Bullock
RLC	Roy L. Caldwell
RHG	Robert H. Gore
RHM	R. H. Millar
RM	Raymond B. Manning
RO	Randy Olson
RS	Rick Steger
SS	Stephen Shuster
SW	Suelynn Williams
VB	Victoria Batista

LITERATURE REFERENCES

B, D, F, G = appendices B, D, F, and G in Birkeland et al., 1976.

E73, E74, E75, E76 = reports of the Environmental Sciences Program of the Smithsonian Institution (Environmental Monitoring and Baseline Data from the Isthmus of Panama - 1973, 1974, 1975, 1976, respectively).

a	= Fauchald (1977)
b	= Birkeland (1974)
c	= Bertsch (1975a)
d	= Reimer (1975)
e	= Gore and Abele (1976)
f	= Macintyre and Glynn (1976)
g	= Sebens (1977)
h	= Meyer (1977)
i	= Sebens (1976)
j	= Bertsch (1975b)
k	= Hendler (1977)
m	= Abrams (1976)
n	= Koehl (1977)
o	= Yee and Chang (1978)
p	= Lawrence (1976)
q	= Sebens (1982)

r = Sebens and DeRiemer (1977)
 s = Gore (1977)
 t = Porter (1972)
 u = Gore and Abele (1973)
 v = Olson (1979)
 w = Birkeland et al. (1976)
 x = Vasquez-Montoya (1979)
 y = Batista (1980)
 z = Abele (1972)
 aa = Powell (1971)
 bb = Smith (1973)
 cc = Meyer (1973)
 dd = Brattgard (1974)
 ee = Lasker (1979)
 ff = Lehman and Porter (1973)
 gg = Spivey (1976)
 hh = Henry and McLaughlin (1975)
 ii = Jackson (1973)
 jj = Rützler and Sterrer (1970)
 kk = Child (1979)
 ll = Southward (1975)
 mm = Caldwell (1981)

OTHER

+ signifies this species is currently represented in the reference collection of the Galeta Point Marine Laboratory.

PORIFERA

<u>Anthosigmella varians</u>	E73, E74, E75, E76, v, q
<u>Craniella</u> sp.	E73, E74, E75, E76
<u>Geodia</u> sp.	F
<u>Niphates erecta</u>	KR, y
<u>Placospongia</u> sp.	E75
<u>Sigmadocia caerulea</u>	KR, y
<u>Sigmadocia</u> sp.	KR, y
<u>Spongia</u> sp.	KR, y
<u>Tedania ignis</u>	KR, y

COELENTERATA

(by Order)

HYDROIDA

<u>Millepora alcicornis</u>	t
<u>Millepora complanata</u>	B, E73, E74, E76, f, q, v, t, jj
<u>Millepora moniliformis</u>	t
<u>Millepora</u> spp.	v
<u>Stylaster rosaceus</u>	DLM, +
Unidentified sp.	y

CERIANTHARIA

<u>Ceriantheopsis americanus</u>	y
<u>Cerianthus</u> sp.	B, +

ACTINIARIA

<u>Aiptasia tagetes</u>	B, q, y, +
<u>Anthopleura krebsi</u>	B, F, +
<u>Anthopleura</u> sp.	B, E73, +
<u>Bartholomea annulata</u>	r, y
<u>Bunodeopsis antilliensis</u>	r
<u>Bunodeopsis globulifera</u>	KS, ? in y
<u>Condylactis gigantea</u>	i, r, bb
<u>Epiphellia</u> n. sp.	B, q, +
<u>Lebrunia coralligens</u>	r
<u>Lebrunia danae</u>	r
<u>Phyllactis flosculifera</u>	q, P. <u>flosculifera</u> [sic] in CC, B, E73, E74, E75, E76, +
<u>Phymanthus crucifer</u>	CC, B, E73, E74, E76, q, r, +
<u>Phymanthus</u> sp.	E75
<u>Stoichactis helianthus</u>	CC, i, r
<u>Telmatactis americana</u>	B, +
<u>Telmatactis roseni</u>	B, +
<u>Telmatactis vernonia</u>	r

CORALLIMORPHARIA

<u>Paradiscosoma neglecta</u>	r
<u>Rhodactis sancti-thomae</u>	r
<u>Ricordea florida</u>	r, v, <u>Recordia</u> in CC, JRY, +

ZOANTHIDEA

<u>Isaurus duchassaingi</u>	AR, B, E73, E76, q, +
<u>Isaurus tuberculatus</u>	q
<u>Palythoa caribaeorum</u>	B, E73, E74, E76, g, n, q, v, +
<u>Palythoa gigantea</u>	E74
<u>Palythoa</u> sp.	E75, E76
<u>Palythoa</u> spp.	v
<u>Palythoa sociatus</u>	y
<u>Palythoa variabilis</u>	B, E74, d, g, n, q, y, +
<u>Palythoa</u> (= <u>Protopalythoa</u>) <u>grandis</u>	q
<u>Parazoanthus parasiticus</u>	q
<u>Parazonanthus swiftii</u>	q
<u>Zoanthus pulchellus</u>	AR, +
<u>Zoanthus sociatus</u>	AR, B, F, E73, E74, E75, E76, g, d, q, y, +
<u>Zoanthus solanderi</u>	AR, B, E73, E74, E75, E76, + <u>Z. solandri</u> [sic] in g, q
<u>Zonanthus</u> sp.	E76

RHIZOSTOMAE

<u>Cassiopea ?xamachana</u>	JC
-----------------------------	----

SCLERACTINIA (by family)

Astrocoeniidae

<u>Stephanocoenia michelinii</u>	JW, f, <u>Stephanocoenia</u> [sic] in t, +
----------------------------------	---

Pocilloporidae

<u>Madracis decatis</u>	JW, t, +
-------------------------	----------

Acroporidae

<u>Acropora palmata</u>	t, v
<u>Acropora cervicornis</u>	t

Agariciidae

<u>Agaricia agaricites</u>	JWO, B, E73, E74, E76, f, v, +
<u>Agaricia agaricites</u> forma <u>agaricites</u>	t
<u>Agaricia agaricites</u> forma <u>danae</u>	t
<u>Agaricia agaricites</u> forma <u>crassa</u>	t
<u>Agaricia agaricites</u> forma <u>purpurea</u>	t
<u>Agaricia agaricites</u> forma <u>humilis</u>	t
<u>Agaricia tenuifolia</u>	t
<u>Agaricia</u> spp.	v
<u>Helioseris cucullata</u>	t

Siderastreidae

<u>Siderastrea siderea</u>	DLM, B, f, q, t, v, y, +
<u>Siderastrea radians</u>	B, E73, E74, f, t, y, jj

Poritidae

<u>Porites astreoides</u>	JW, B, E73, E74, E76, f, t, v, y, + <u>P. astreoides</u> in E75, jj
<u>Porites branneri</u>	JWP, t, +
<u>Porites divaricata</u>	t
<u>Porites furcata</u>	B, E73, E74, E76, f, t, jj
<u>Porites porites</u>	t, y
<u>Porites</u> sp.	E76

Faviidae

<u>Colpophyllia amaranthus</u>	t
<u>Colpophyllia breviserialis</u>	t
<u>Colpophyllia natans</u>	JW, t, +
<u>Diplora clivosa</u>	t, q, v
<u>Diploria strigosa</u>	t
<u>Favia fragum</u>	DLM, B, E73, E74, E76, f, t, y, +
<u>Manicina areolata</u>	JWP, t, +
<u>Montastrea annularis</u>	JW, f, t, +
<u>Montastrea cavernosa</u>	JW, t, +, ee, ff

Rhizangiidae

Astrangia solitaria B, t

Meandrinidae

<u>Dichocoenia cf. stellaris</u>	JWP, +
<u>Dichocoenia stokesii</u>	B, f, t
	<u>D. stokesia</u> in E76
<u>Meandrina meandrites</u>	t
<u>Meandrina meandrites</u>	
var. <u>meandrites</u>	JW, +

Mussidae

<u>Isophyllia multiflora</u>	t
<u>Isophyllia sinuosa</u>	PG, f, t, +
<u>Mussa angulosa</u>	JWP, t, +
<u>Mycetophyllia lamarckana</u>	JW, t, +
<u>Mycetophyllia</u> sp. B	t

Caryophyllidae

Eusmilia fastigiata t

Dendrophylliidae

Tubastraea aurea t

GORGONACEA

<u>Erythropodium caribaeorum</u>	B, E73, E74, E76, q
<u>Gorgonia flabellum</u>	f
<u>Gorgonia</u> spp.	v
<u>Gorgonia ventalina</u>	b

PLATYHELMINTHES

Class: Turbellaria

Unident. species y

SIPUNCULA

<u>Aspidosiphon broki</u>	B, +
<u>Aspidosiphon speculator</u>	x
<u>Aspidosiphon spinoscutatus</u>	x
<u>Aspidosiphon</u> spp. (7 species)	B
<u>Dendrostomum</u> sp.	B
<u>Golfingia rimicola</u>	x
<u>Golfingia</u> spp. (4 species)	B, F
<u>Lithacrosiphon</u> spp. (5 species)	B
<u>Lithacrosiphon</u> sp. 3	F
<u>Paraspidosiphon fisheri</u>	B, F, +
<u>Paraspidosiphon speciosus</u>	? in B
<u>Paraspidosiphon spinoso-scutatus</u>	B, F, +
<u>Paraspidosiphon steenstrupi</u>	B, F, +
<u>Paraspidosiphon</u> spp. (7 species)	B
<u>Paraspidosiphon</u> sp. 4	F
<u>Phascolosoma antillarum</u>	B, F, +
<u>Phascolosoma perlucens</u>	B, F, +
<u>Phascolosoma varians</u>	B, +
<u>Phascolosoma</u> spp. (7 species)	B
<u>Phascolosoma</u> spp. 3, 4	F
<u>Themiste</u> spp. (3 species)	B
Unident. species	y

ANNELIDA

(by family)

Ampharetidae

<u>Isolda bipinnata</u>	a
<u>Isolda pulchella</u>	x
<u>Melinna</u> n. sp.	B

Amphinomidae

<i>Amphinomid</i> sp. 1	B, F, +
<i>Amphinomid</i> sp. 2	B, +
<u>Eurythoe complanata</u>	B, F, a, q, +
<u>Hermodice carunculata</u>	B, a, q, y
<u>Linopherus canariensis</u>	a (family Phyllodocidae in B)
Unident. species	y

Aphroditidae

<u>Aphrodita diplops</u>	a
<u>Aphrodita</u> n. sp.	B

Arabellidae

<u>Arabella mutans</u>	B, F, a, x, y, +
<u>Arabella</u> sp. indet.	a

Ariciidae (listed thus by Vasquez-Montoya, 1979)

<u>Naineris laevigata</u>	x
<u>Naineris mutila</u>	x
<u>Naineris setosa</u>	x
<u>Scoloplos armiger</u>	x

Chrysopetalidae (Palmyridae in B)

<u>Bhawania goodei</u>	B, a, +
<u>Bhawania riveti</u>	+

Cirratulidae

<u>Caulieriella hamata</u>	x
<u>Caulieriella</u> sp. indet.	B, a
<u>Chaetozone</u> sp. indet.	B, a, y
<u>Cirratulus cirratus</u>	? in a, <u>C. cirratulus</u> in B
<u>Cirriformia luxuriosa</u>	B, + (specimen in collection from Pacific)
<u>Cirriformia punctata</u>	B, a, +

<u>Dodecaceria concharum</u>	B, E74, a
<u>Tharyx</u> sp. indet.	B, a
 Dorvilleidae	
<u>Dorvillea rubrovittatus</u>	a, <u>D. rubrovittata</u> in B
<u>Driloneresis nuda</u>	x
<u>Schistomerengos longicornis</u>	a, x
 Eunicidae (Leodicidae in F)	
<u>Eunice afra</u>	B, F, a, +
<u>Eunice antennata aedificatrix</u>	B, a, +
<u>Eunice aphroditoides</u>	B, F, a, +
<u>Eunice (Nicidion) caribaea</u>	a, +, <u>E. caribaea</u> [sic] in B, F
<u>Eunice filamentosa</u>	B, a, +
<u>Eunice vittatopsis</u>	x
<u>Eunice websteri</u>	B, a, +
<u>Eunice</u> sp. indet.	B, a
<u>Lysidice ninetta</u>	x
<u>Lysidice</u> sp.	y
<u>Marpheya amadae</u>	a
<u>Marpheya depressa</u>	x
<u>Marpheya</u> n. sp.	B
<u>Marpheya</u> sp. indet.	a
<u>Nematonereis unicornis</u>	a, x
<u>Palola siciliensis</u>	B, a, +
<u>Palola</u> sp. indet.	B, a
 Flabelligeridae (Chloraemidae in B)	
<u>Pherusa capulata</u>	x
<u>Pherusa inflata</u>	B, a, +
<u>Piromis americana</u>	+ (no identifier on voucher specimen)
 Glyceridae	
<u>Glycera abranchiata</u>	x

<u>Glycera oxycephala</u>	B, F, a
<u>Glycera tesselata</u>	B, a
<u>Glycera</u> sp.	B

Goniadidae

<u>Goniada acicula</u>	B, a
------------------------	------

Hesionidae

<u>Hesione picta</u>	B, a
<u>Ophiodromus obscurus</u>	B, a

Lumbrineridae

<u>Lumbrineris inflata</u>	a, <u>Lumbrinereis</u> in B
<u>Lumbrineris</u> sp. aff. <u>latreilli</u>	y
<u>Lumbrineris tetraura</u>	x, y

Lysaretidae

<u>Lysaretid</u> sp.	B, a
<u>Oenone fulgida</u>	B, a (in family Arabellidae in B)

Maldanidae

<u>Axiothella rubrocincta</u>	x
-------------------------------	---

Nereidae

<u>Ceratonereis mirabilis</u>	B, a, +
<u>Neanthes galetae</u>	a, x
<u>Neanthes</u> n. sp. 1	B
<u>Neanthes</u> sp. indet.	B, a
<u>Nematonereis</u> sp.	y
<u>Nereis callaona</u>	B, F, a, +
<u>Nereis panamensis</u>	a
<u>Nereis</u> n. sp. A	B
<u>Nereis riisei</u>	a, <u>riseii</u> [sic] in B, x
<u>Nereis</u> sp.	y
<u>Nereidae</u> sp. indet.	a

<u>Perinereis elenacasoii</u>	B, F, +
<u>Perinereis anderssoni</u>	a, x, y
<u>Perinereis</u> sp. indet. A	B
<u>Perinereis</u> sp. indet. B	B
<u>Platynereis dumerilii</u>	B, F, a, x, +
<u>Platynereis</u> sp. indet.	B, a
<u>Pseudonereis gallapagensis</u>	B, F, +
Onuphidae	
<u>Onuphis nebulosa</u>	x
<u>Onuphis vermillionensis</u>	B, a, x (in family Eunacidae in B)
<u>Onuphis</u> sp.	B (in family Eunacidae in B)
Opheliidae	
<u>Armandia bioculata</u>	x
Oweniidae	
<u>Owenia collaris</u>	B, a
Paraonidae	
<u>Aricidea suecica</u>	x
Phyllodocidae	
<u>Anaitides erythrophylloides</u>	a, <u>erythrophylla</u> [sic] in B
<u>Anaitides</u> sp. cf. <u>lamellifera</u>	x
<u>Eulalia myriacyclum</u>	B, F, a, +
<u>Sige orientalis</u>	B, ? in a
Poecilochaetidae	
<u>Poecilochaetus johnsoni</u>	x
Polynoidae	
<u>Halosydna leucohyba</u>	B, a, +
<u>Halosydna</u> sp. 1	B, +

<u>Harmothoe</u> sp. indet.	B, a
<u>Harmothoe</u> <u>hirsuta</u>	x
<u>Lepidonotus</u> <u>humilis</u>	B, a, +
<u>Lepidonotus</u> <u>neophilus</u>	+
<u>Lepidasthenia</u> <u>varius</u>	a

Sabellariidae

<u>Phragmatopoma</u> sp. indet.	B, a
<u>Sabellaria</u> <u>alcocki</u>	B, + (specimen in collection from Pacific)
<u>Sabellaria</u> <u>floridensis</u>	B, a, +

Sabellidae

<u>Chone</u> sp.	B, a
<u>Demonax</u> <u>leucaspis</u>	? in a
<u>Demonax</u> sp.	B
<u>Hypsicomus</u> <u>torquatus</u>	+, B, F
<u>Hypsicomus</u> <u>phaeotenia</u>	a
<u>Megalomma</u> sp.	y
<u>Megalomma</u> sp. aff. <u>pigmentum</u>	x
<u>Megalomma</u> <u>roulei</u>	x
<u>Megalomma</u> <u>vesiculosum</u>	a, <u>Megaloma</u> [sic] in B
<u>Pseudopotamilla</u> <u>reniformis</u>	B, a
<u>Sabella</u> <u>melanostigma</u>	B, a, +
<u>Sabella</u> sp. 17	B
<u>Sabella</u> sp. 18	B
<u>Sabella</u> sp.	y
Sabellidae sp. indet.	a
<u>Sabellastarte</u> <u>magnifica</u>	JC

Serpulidae

Serpulid sp. 1	F
Serpulid sp. 2	F
<u>Spirobranchus</u> <u>giganteus</u>	JC
<u>Spirorbis</u> sp.	F, y

Sigalionidae

<u>Psammolyce spinosa</u>	B, a, +
<u>Sthenelais verruculosa</u>	x

Spionidae

<u>Boccardia polybranchia</u>	x
<u>Malacoceros indicus</u>	x
<u>Nerinides cantabra</u>	x
<u>Prionospio cirrifera</u>	x
<u>Prionospio heterobranchia</u>	
<u>texana</u>	B, a
<u>Pseudopolydora antennata</u>	x

Syllidae

<u>Autolytus anoplos</u>	a
<u>Autolytus</u> n. sp.	B
<u>Autolytus</u> cf. <u>magnus</u>	B, F, +
<u>Haplosyllis spongicola</u>	B, a, +
<u>Langerhansia cornuta</u>	B, B, a, +
<u>Langerhansia mexicana</u>	B, a
<u>Odontosyllis</u> sp.	B
<u>Opisthosyllis brunnea</u>	B, F, a, +
<u>Pionosyllis</u> sp. indet.	a
Syllidae, unident. fragments	a
<u>Syllis longissima</u>	x
<u>Trypanosyllis taeniaformis</u>	x, y
<u>Typosyllis aciculata</u>	B, a, + (specimen in collection from Pacific)
<u>Typosyllis fuscosuturata</u>	a
<u>Typosyllis</u> sp. A	B
<u>Typosyllis prolifera</u>	B, ? in a
<u>Typosyllis variegata</u>	B, a, +

Terebellidae

<u>Eupolynnia nebulosa</u>	B, F, a, +
----------------------------	------------

<u>Euthelepus pascua</u>	a
<u>Loimia medusa</u>	B, F, a
<u>Pista fasciata</u>	B, a, +
<u>Polycirrus</u> sp.	B, a, x
<u>Polycirrus</u> sp. aff. <u>haematodes</u>	x
<u>Streblosoma crassibranchia</u>	B, a, +
Terebellidae sp. indet.	a
<u>Thelepus setosus</u>	B, a

Trichobranchidae

<u>Trebellides stroemi</u>	x
----------------------------	---

ARTHROPODA

CRUSTACEA: COPEPODA (by order)

CALANOIDA

Unident. species	y
------------------	---

CRUSTACEA: ISOPODA (by family)

Anthuridae

<u>Accalathura</u> sp.?	ss
-------------------------	----

Idoteidae

<u>Cleantis planicauda</u>	x
----------------------------	---

Cirolanidae

<u>Cirolana parva</u>	ss
<u>Excirolana mayana</u>	x

Excorallanidae

<u>Alcinora</u> sp.	y
<u>Excorallana tricornis</u>	ss

Ligiidae

<u>Ligia</u> sp.	y
------------------	---

Limnoriidae

<u>Limnoria</u> sp.	y
---------------------	---

Sphaeromatidae

<u>Paracerceis caudata</u>	SS
----------------------------	----

Stenetriidae

<u>Stenetrium serratum</u>	SS
----------------------------	----

CRUSTACEA: AMPHIPODA (by family)

Gammaridae

<u>Elasmopus</u> sp.	y
----------------------	---

CRUSTACEA: DECAPODA: REPTANTIA (by family)

Porcellanidae

<u>Clastotoechus nodosus</u>	B, JH, e
<u>Megalobrachium poeyi</u>	B, JH, e, +
<u>Megalobrachium roseum</u>	B, JH, e, +
<u>Megalobrachium soriatum</u>	B, JH, e, +
<u>Neopisosoma angustifrons</u>	e, s, +
<u>Pachycheles chacei</u>	B, JH, e, +
<u>Pachycheles cristobalensis</u>	B, JH, e, +
<u>Pachycheles serratus</u>	B, JH, JS, LGA, e, +
<u>Pachycheles susanae</u>	B, e, u, +
<u>Petrolisthes armatus</u>	B, JH, e, +
<u>Petrolisthes galathinus</u>	B, JH, JS, e, +
<u>Petrolisthes jugosus</u>	B, JH, e, +

Grapsidae

<u>Aratus pisonii</u>	EK, y
-----------------------	-------

<u>Goniopsis cruentata</u>	RHG, y
<u>Grapsus grapsus</u>	RHG, +
<u>Pachygrapsus gracilis</u>	y
<u>Pachygrapsus marmoratus</u>	B
<u>Pachygrapsus transversus</u>	B, JSG, +
<u>Percnon gibbesi</u>	RHG
<u>Plagusia depressa</u>	RHG, JPS, +
<u>Sesarma cinereum</u>	y
<u>Sesarma curacaoense</u>	y

Portunidae

<u>Callinectes</u> sp.	JC
------------------------	----

Majidae

<u>Acanthonyx petiverii</u>	JSG, + <u>petriverii</u> [sic] in B
<u>Epialtus</u> sp.	B
= <u>E. ?longirostris</u>	JSG
<u>Macrocoeloma subparallelum</u>	B, JSG
<u>Microphrys bicornutus</u>	B, F, JSG, E74, E75, E76, x, y, +
<u>Mithrax acuticornis</u>	B, JSG
<u>Mithrax coryphe</u>	B, JSG, +
<u>Mithrax commensalis</u>	JRS, JS
<u>Mithrax sculptus?</u>	EK
<u>Mithrax spinosissimus</u>	B
<u>Mithrax</u> sp.	B, JSG
<u>Mithrax verrucosus</u>	JRS, JS
<u>Pitho aculeata</u>	B, JSG, +
<u>Stenorhynchus seticornis</u>	JC
<u>Thoe puella</u>	B, JSG

Xanthidae

<u>Carpilius corallinus</u>	JC
<u>Cataleptodius floridanus</u>	RHG, <u>Leptodium</u> in B, JSG, LGA, +
<u>Domecia acanthophora</u>	B, JSG
<u>Eriphia gonagra</u>	B, JSG

<u>Leptodius floridanus</u>	x
<u>Micropanope</u> sp.	B, JSG
<u>Panopeus bermudensis</u>	B, JSG, LGA
<u>Panopeus harttii</u>	RHG, +
<u>Panopeus herbstii</u>	JSG, x, y, +
<u>Panopeus</u> sp.	B, JSG
<u>Panopeus</u> sp. aff. <u>occidentalis</u>	x
<u>Paraliomera dispar</u>	B, JSG, LGA, +
<u>Pilumnus dasypodus</u>	B, JSG, +
<u>Pilumnus holosericus</u>	B, JSG, LGA, +
<u>Pilumnus lacteus</u>	B
<u>Pilumnus reticulatus</u>	B, JSG, +
<u>Pilumnus sayi?</u>	RHG, +
<u>Pilumnus</u> sp.	B, JSG
<u>Platyactaea setigera</u>	RHG, <u>Actaea</u> in B, JSG, +
<u>Platypodia spectabilis</u>	B, JSG, LGA, RHG, +
<u>Xantho denticulatus</u>	RHG, <u>Xanthodius</u> in B, JSG, +

Calappidae

<u>Calappa</u> sp.	JC
--------------------	----

Gecarcinidae

<u>Cardisoma guanhumi</u>	RHG
---------------------------	-----

Dromiidae

<u>Dromidia</u> sp.	JC
---------------------	----

Ocypodidae

<u>Ocypode quadrata</u>	y
<u>Uca burgersi</u>	RHG, +
<u>Uca</u> sp.	y, jj

Palinuridae

<u>Panulirus argus</u>	JC
<u>Panulirus guttatus</u>	JC

Scyllaridae

Scyllarus sp. JC

Parthenopidae

Heterocrypta macrobranchia? B, RHG

Leucosiidae

Uhlias limbatus B, JSG

Paguridae

<u>Paguristes cadenati</u>	JC
<u>Paguristes grayi</u>	PAA, m, +
<u>Paguristes cf tortugae</u>	PAA, m, +
<u>Pagurus bonairensis</u>	PAA, m, +
<u>Pagurus brevidactylus</u>	PAA, m, +

Coenobitidae

Coenobita clypeatus PAA, m, +

Diogenidae

<u>Calcinus tibicen</u>	PAA, JH, m, +
<u>Clibanarius antillensis</u>	PAA, m, +
<u>Clibanarius</u> sp.	y
<u>Clibanarius tricolor</u>	PAA, m, +
<u>Dardanus venosus</u>	PAA, m, +
<u>Dardanus</u> sp.	o
<u>Petrochirus diogenes</u>	RHG, <u>P. bahamensis</u> in PAA, m, +
<u>Petrochirus</u> sp.	o

Pinnotheridae

<u>Pinnotheres maculatus</u>	JSG
<u>Pinnixa ?faxonii</u>	JSG

Hapalocarcinidae

Pseudocryptochirus RHG

CRUSTACEA: DECAPODA: NATANTIA (by family)

Alpheidae

<u>Alpheus armillatus</u>	B, F, x
<u>Alpheus armatus</u>	JC
<u>Alpheus bahamensis</u>	B
<u>Alpheus cristulifrons</u>	B
<u>Alpheus floridanus</u>	x
<u>Alpheus formosus</u>	B
<u>Alpheus normanni</u>	B
<u>Alpheus nuttingi</u>	B
<u>Alpheus paracrinitus</u>	B
<u>Alpheus peasei</u>	B
<u>Alpheus ridleyi</u>	B
<u>Alpheus schmitti</u>	B
<u>Alpheus simus</u>	B
<u>Alpheus sp.</u>	B, F, y
<u>Alpheus viridari</u>	F, x
<u>Automate rectifrons</u>	B
<u>Metalpheus rostratipes</u>	B
<u>Salmoneus ortmanni</u>	B
<u>Synalpheus anasimus</u>	B
<u>Synalpheus brevidactylus</u>	RHG
<u>Synalpheus fritzmuelleri</u>	B
<u>Synalpheus herricki</u>	B
<u>Synalpheus minus</u>	
<u>Synalpheus pandionis</u>	B
<u>Synalpheus sp.</u>	B
<u>Synalpheus tenuispina</u>	B
<u>Synalpheus townsendi</u>	B
<u>Thunor rathbunae</u>	B

Callianassidae

<u>Callichirus acanthochinus</u>	x
----------------------------------	---

Gnathophyllidae

<u>Gnathophyllum americanum</u>	B
---------------------------------	---

Hippolytidae

<u>Hippolyte curacaoensis</u>	B
<u>Lysmata intermedia</u>	B
<u>Thor manningi</u>	B, x

Palaemonidae

<u>Periclimenes americanus</u>	B
<u>Stenopus hispidus</u>	y
<u>Stenopus scutellatus</u>	JC

Penaeidae

<u>Penaeus duorarum</u>	x
<u>Penaeus</u> sp.	mm
<u>Sicyonia parri</u>	B
<u>Trachypenaeus similis</u>	B

Processidae

<u>Ambidexter symmetricus</u>	B, B, x, z
<u>Processa bermudensis</u>	B
<u>Processa fimbriata</u>	B
<u>Processa</u> sp. aff. <u>hemphilli</u>	x

CRUSTACEA: STOMATOPODA

<u>Gonodactylus austrinus</u>	RM, B, y, +
<u>Gonodactylus bredini</u>	RM, B, y, +
<u>Gonodactylus oerstedi</u>	RM, B, y, +
<u>Gonodactylus spinulosus</u>	HD and RS, +
<u>Meiosquilla lebouri</u>	RLC and RS
<u>Pseudosquilla ciliata</u>	RM, x, y, +
<u>Nannosquilla</u> sp.	x

CRUSTACEA: MYSIDACEA

<u>Siriella chierchiae</u>	dd
<u>Bowmaniella bracescui</u>	dd
<u>Bowmaniella sewelli</u>	dd
<u>Amathimysis cherados</u>	dd
<u>Amathimysis gibba</u>	dd
<u>Brasiliomysis castroi</u>	dd
<u>Cubanomysis jimenezi</u>	dd
<u>Mysidopsis brattstroemi</u>	dd
<u>Mysidopsis velifera</u>	dd
<u>Mysidopsis arenosa</u> sp. nov.	dd
<u>Mysidium columbiae</u>	dd
<u>Mysidium gracile</u>	dd
<u>Mysidium integrum</u>	dd

CRUSTACEA: CIRRIPEDIA

<u>Balanus</u> sp.	jj
<u>Balanus trigonus</u>	gg
<u>Balanus venustus</u>	hh
<u>Chthamalus angustitergum</u>	ll
<u>Chthamalus bisinuatus</u>	ll
<u>Chthamalus rhizophorae</u>	ll
<u>Chthamalus</u> sp.	y
<u>Newmanella radiata</u>	gg
<u>Tetraclita stalactifera</u>	ll

PYCNOGONIDA

<u>Achelia sawayai</u>	B, kk, +
<u>Anoplodactylus allotrius</u>	kk
<u>Anoplodactylus batangensis</u>	<u>batangense</u> in B, kk
<u>Anoplodactylus evelinae</u>	B, kk, + (specimen in collection from Pacific)

<u>Anoplodactylus galetensis</u>	kk
<u>Anoplodactylus insigniformis</u>	kk
<u>Anoplodactylus jonesi</u>	kk
<u>Anoplodactylus monotrema</u>	kk
<u>Anoplodactylus multiclavus</u>	kk
<u>Anoplodactylus pectinus</u>	kk
<u>Anoplodactylus</u> spp.	B, kk
<u>Anoplodactylus stri</u>	kk
<u>Anoplodactylus trispinosus</u>	kk
<u>Anoplodactylus viridintestinalis</u>	kk
<u>Ammothella appendiculata</u>	B, kk, +
<u>Ammothella exornata</u>	kk, +
<u>Ammothella marcusi</u>	kk
<u>Ammothella rugulosa</u>	kk
<u>Ammothella</u> spp.	B
<u>Ammothella spinifera</u>	kk
<u>Ascorhynchus castellioides</u>	B, kk, +
<u>Ascorhynchus latipes</u>	B, kk, +
<u>Callipallene emaciata</u>	kk
<u>Callipallene</u> sp.	B
<u>Endeis spinosa</u>	kk
<u>Eurycyde gorda</u>	kk
<u>Eurycyde raphiaster</u>	B, kk, +
<u>Eurycyde</u> sp.	B
<u>Nymphon floridanum</u>	CAC, kk, +
<u>Nymphopsis duodorsospinosa</u>	B, kk, + (specimen in collection from Pacific)
<u>Pallenopsis schmitti</u>	kk
<u>Pigrogromitus timsanus</u>	B, kk, +
<u>Rhynchothorax architectus</u>	kk
<u>Rhynchothorax</u> sp.	B
<u>Tanystylum birkelandi</u>	kk
<u>Tanystylum geminum</u>	kk
<u>Tanystylum isthmiacum</u>	
<u>difficile</u>	kk
<u>Tanystylum</u> sp.	B, kk

MOLLUSCA

SCAPHOPODA

<u>Dentalium gouldi</u>	KBM, +
<u>Dentalium antillarum</u>	x

POLYPLACOPHORA

<u>Acanthochitona hemphilli</u>	B
<u>Acanthochitona interfissa</u>	B
<u>Acanthochitona pygmaea</u>	B
<u>Acanthochitona spiculosus</u>	B
<u>Acanthochitona unident. sp.</u>	y
<u>Acanthopleura granulata</u>	B
<u>Calloplax janeirensis</u>	B
<u>Chiton viridis</u>	B
<u>Choneplax lata</u>	B
<u>Ischnochiton papillosum</u>	B
<u>Ischnochiton pectinatus</u>	B
<u>Ischnochiton purpurascens</u>	B
<u>Lepidochitona liozonis</u>	B

GASTROPODA: PROSOBRANCHIA

<u>Acmaea antillarum</u>	B
<u>Acmaea pustulata</u>	B, f
<u>Alvania aberrans</u>	CB, +
<u>Anachis catenata</u>	CB, B, +
<u>Anachis crassilabris</u>	F
<u>Anachis obesa</u>	JR, CB, +
<u>Antillophos</u> sp. 1	CB, +
<u>Architectonica nobilis</u>	KBM, +
<u>Arene cruentata</u>	CB, B, +
<u>Arene tricarinata</u>	B

<u>Astraea caelata</u>	DLW, B, +
<u>Astraea phoebia</u>	KBM, IT, B, +
<u>Bailya intricata</u>	B, F
<u>Bailya parva</u>	KBM, +
<u>Balcis intermedia</u>	B
<u>Balcis</u> sp. 1	B
<u>Batillaria minima</u>	KBM, D, F, f, y, +
<u>Bittium varium</u>	B
<u>Bursa cubaniana</u>	B
<u>Bursa granularis</u>	KBM, +
<u>Caecum</u> sp.	y
<u>Calliostoma jujubinum</u>	KBM, +
<u>Cantharus auritulus</u>	F
<u>Cantharus tinctus</u>	B
<u>Cerithiopsis emersoni</u>	F
<u>Cerithium eburneum</u>	KBM, B, f, x, +
<u>Cerithium litteratum</u>	KBM, B, JR, f, x, y, + [cf. <u>C litteratum</u> (Young) by CB in +]
<u>Cerithium variabile</u>	B, D
<u>Charonia variegata</u>	B, o, +
<u>Cheilea equestris</u>	B, <u>Cheila</u> [sic] in B
<u>Cittarium pica</u>	B
<u>Columbella mercatoria</u>	KBM, +
<u>Conus daucus</u>	o
<u>Conus mus</u>	B
<u>Coralliophila abbreviata</u>	KBM, +
<u>Coralliophila aberrans</u>	CB, +
<u>Coralliophila caribaea</u>	B
<u>Crassispira fuscescens</u>	KBM, +
<u>Crassispira leucocyma</u>	CB, +
<u>Crassispira nigrescens</u>	JR
<u>Crassispira tampaensis</u>	x
<u>Crepidula plana</u>	B
<u>Crucibulum auricula</u>	KBM, f, +
<u>Cyclostremiscus beauvi</u>	B

<u>Cymatium muricinum</u>	KBM, +
<u>Cymatium nicobaricum</u>	KBM, B, +
<u>Cymatium pileare</u>	KBM, B, o, +
<u>Cyphoma gibbosum</u>	KBM, b, +
<u>Cyprea cinerea</u>	KBM, +
<u>Cyprea zebra</u>	KBM, B, o, +
<u>Cypraeocassis testiculus</u>	KBM, B, E76, k, o, +
<u>Daphnella lymneiformis</u>	CB, B, +
<u>Diodora cayenensis</u>	CB, B, +
<u>Diodora dysoni</u>	CB, B, JR, +
<u>Diodora cf. minuta</u>	KBM, B, JR, +
<u>Diodora variegata</u>	B
<u>Drillia albinodata</u>	RS
<u>Drillia</u> sp. 1	CB, JR, +
<u>Drupa nodulosa</u>	B, D, <u>Morula nodulosa</u> KBM, +
<u>Emarginula phrixodes</u>	B
<u>Emarginula pumila</u>	B
<u>Engina turbinella</u>	CB, +
<u>Epitonium candeanum</u>	B
<u>Epitonium lamellosum</u>	B
<u>Epitonium occidentale</u>	B
<u>Epitonium</u> sp. 1	B
<u>Fasciolaria tulipa</u>	KBM, B, o, x, +
<u>Fissurella angusta</u>	B
<u>Fissurella barbadensis</u>	B
<u>Heliacus bisulcatus</u>	B
<u>Heliacus cylindricus</u>	B
<u>Heliacus infundibuliformis</u>	B, <u>infunibuliformis</u> [sic] in F
<u>Hemitoma octoradiata</u>	JR, DLW, +
<u>Hemitoma</u> sp.	B
<u>Hipponix antiquatus</u>	CB, B, f, +
<u>Hipponix subrufus</u>	B
<u>Hyalina albolineata</u>	B
<u>Hyalina avena</u>	KBM, B, F, x, +
<u>Hyalina tenuilabra</u>	B

<u>Janthina janthina</u>	KBM, +
<u>Latirus carniferus</u>	RB, KBM, JR, +, <u>carnifera</u> [sic] in B
<u>Latirus distinctus</u>	y
<u>Latirus infundibulum</u>	o
<u>Leucozonia nassa</u>	RB, +
<u>Leucozonia ocellata</u>	KBM, B, +
<u>Littorina angulifera</u>	KBM, y, +
<u>Littorina angustior</u>	JC, (KBM, + as <u>L. lineata</u>)
<u>Littorina lineolata</u>	B, D
<u>Littorina meleagris</u>	D
<u>Littorina nebulosa</u>	KBM, y, +
<u>Littorina tessellata</u>	KBM, +
<u>Littorina ziczac</u>	KBM, D, +
<u>Lucapina suffusa</u>	KBM, +
<u>Mangelia fusca</u>	B, +
<u>Marginella gracilis</u>	x
<u>Melongena melongena</u>	KBM, x, +
Melogenidae sp.	o
<u>Mitra nodulosa</u>	KBM, +
<u>Mitra</u> sp.	B
<u>Modulus modulus</u>	KBM, B, f, +
<u>Morula nodulosa</u>	see <u>Drupa nodulosa</u>
<u>Morum oniscus</u>	KBM, o, +
<u>Murex dilectus</u>	KBM, +
<u>Murex pomum</u>	KBM, KBM, o, +
<u>Murex recurvirostris rubidus</u>	KBM, G, GLH, +
cf. <u>Murex woodringi</u>	KBM, +
<u>Nassarius vibex</u>	KBM, F, x, +
<u>Neosimnia acicularis</u>	CB, +
<u>Nerita fulgurans</u>	KBM, D, +
<u>Nerita peloronta</u>	KBM, D, +
<u>Nerita tessellata</u>	KBM, D, y, +
<u>Nerita versicolor</u>	KBM, D, +
<u>Neritina virginea</u>	DLW, KBM, JR, f, x, y, +, <u>virgines</u> [sic] in F, <u>virginica</u> [sic] in f

<u>Nitidella nitida</u>	B
<u>Nitidella</u> sp. 2	B
<u>Nodilittorina tuberculata</u>	KBM, D, +
<u>Oliva reticularis</u>	KBM, +
<u>Olivella petiolata</u>	DLW, +
<u>Opalia crenata</u>	CB, B, +
<u>Opalia pumilio</u>	B, <u>pumilo</u> [sic] in B
<u>Petaloconchus</u> sp.	
cf. <u>erectus</u>	y
<u>Pisania pusio</u>	KBM, +
<u>Planaxis lineatus</u>	CB, D, F, +
<u>Planaxis nucleus</u>	D, F
<u>Polinices hepaticus</u>	o, x
<u>Polinices lacteus</u>	KBM, B, + <u>lactens</u> [sic] in f
<u>Prunum guttatum</u>	KBM, JR, +
<u>Psarostola monilifera</u>	B
<u>Purpura patula</u>	KBM, B, o, +
<u>Risomurex muricoides</u>	KBM, B, JR, +
<u>Risomurex roseus</u>	JR, +
<u>Risomurex</u> sp.	JR, CB, +
<u>Rissoina bryerea</u>	B
<u>Rissoina decussata</u>	KBM, B, F, f, +
<u>Rissoina multicostata</u>	KBM, +
<u>Smaragdia viridis</u>	+, <u>S. viridis</u> in KBM, B
<u>Strombus gigas</u>	x
<u>Strombus pugilis</u>	JRY, f, +
<u>Strombus raninus</u>	B, f, x, +
<u>Tectarius muricatus</u>	KBM, D, +
<u>Tegula fasciata</u>	KBM, B, JR, f, +
<u>Thais deltoidea</u>	B, JR, o, +
<u>Thais haemastoma</u> , var A	CB, B, D, JR, +
<u>Thais haemastoma</u>	o
<u>Thais rustica</u>	o
<u>Tonna maculosa</u>	KBM, +
<u>Tricolia adamsi</u>	DLW, KBM, B, +

<u>Tricolia bella</u>	KBM, CB, B, F, +
<u>Tricolia thalassicola</u>	KBM, +
<u>Trivia quadripunctata</u>	DLW, KBM, JR, +
<u>Turritella exoleta</u>	KBM, +
<u>Vasum muricatum</u>	DLW, B, +
<u>Vermidae</u>	y
<u>Voluta musica</u>	DLM, +

GASTROPODA: OPISTHOBRANCHIA (by order)

Anaspidea

<u>Aplysia dactylomela</u>	EK, KBM
<u>Aplysia parvula</u>	KBM
<u>Aplysia</u> sp.	KBM
<u>Bursatella leachii pleii</u>	EK, y, KBM
<u>Dolabridifera dolabridifera</u>	KBM
<u>Petalifera ramosa</u>	KBM
<u>Phyllaplysia engeli</u>	KBM
<u>Stylocheilus longicauda</u>	KBM

Cephalaspidea

<u>Aglaja evelinae</u>	KBM
<u>Atys riiseana</u>	x
<u>Bulla occidentalis</u>	KBM, x
<u>Bulla striata</u>	KBM, f, +
<u>Chelidonura</u> spp.	KBM
<u>Haminoea antillarum</u>	x
<u>Haminoea elegans</u>	KBM
<u>Ildica</u> sp.	KBM
<u>Micromelo undata</u>	DS, SW, KBM

Sacoglossa

<u>Bosellia marcusii</u>	KBM
<u>Bosellia memetica</u>	KBM
<u>Caliphylla mediterranea</u>	KBM

<u>Cyerce antillarum</u>	KBM
<u>Elysia ornata</u>	KBM
<u>Elysia papillosa</u>	KBM
<u>Elysia picta</u>	KBM
<u>Elysia tuca</u>	KBM
<u>Lobiger souverbii</u>	KBM
<u>Oxynoe antillarum</u>	KBM
<u>Placida dendritica</u>	KBM
<u>Phyllobranchillus</u> sp.	KBM
<u>Stiliger</u> sp.	KBM
<u>Tridachia crispata</u>	KBM, +

Notaspidea

<u>Berthella tupala</u>	c
<u>Berthellina quadridens</u>	KBM
<u>Pleurobranchus areolatus</u>	v, KBM

Nudibranchia

<u>Aegires sublaevis</u>	h, KBM
<u>Aphelodoris antillensis</u>	h
<u>Berghia coerulescens</u>	KBM
<u>Berghia creutzbergi</u>	KBM
<u>Bornella calcarata</u>	KBM
<u>Cadlina rumia</u>	j
<u>Catriona tina</u>	KBM
<u>Chromodoris clenchi</u>	h, KBM
<u>Chromodoris kempfi</u>	h, KBM
<u>Chromodoris</u> sp.	y
<u>Coryphella dushia</u>	KBM
<u>Dendrodoris krebsii</u>	KBM
<u>Discodoris evelinae</u>	h
<u>Discodoris mortenseni</u>	j
<u>Dondice occidentalis</u>	KBM

<u>Doriopsilla nigrolineata</u>	h
<u>Doto divae</u>	KBM
<u>Felimare bayeri</u>	h, KBM
<u>Glaucus atlanticus</u>	KBM
<u>Godiva rubolineata</u>	KBM
<u>Hexabranchus sanguineus</u>	h, KBM
<u>Hypselodoris ruthae</u>	h, KBM
<u>Okenia evelinae</u>	h, KBM
<u>Phidiana lynceus</u>	KBM
<u>Phyllidiopsis molaensis</u>	h
<u>Platydoris angustipes</u>	h
<u>Scyllaea pelagica</u>	KBM
<u>Spurilla neopolitana</u>	KBM
<u>Tambja oliva</u>	h, KBM
<u>Tritonia bayeri</u>	KBM
<u>Tritonia wellsi</u>	KBM

GASTROPODA: PULMONATA

<u>Melampus coffeus</u>	KBM, +
-------------------------	--------

BIVALVIA

<u>Americardia media</u>	x
<u>Anodontia pectinata</u>	x
<u>Anomia simplex</u>	DLW, JR, +
<u>Arca imbricata</u>	IT, KBM, B, x, +
<u>Arcopsis adamsi</u>	IT, B, F, f, x, +
<u>Asaphis deflorata</u>	KBM, +
<u>Barbatia domingensis</u>	CB, B, +
<u>Barbatia tenera</u>	BM, B, +
<u>Brachidontes modiolus</u>	JR
<u>Brachidontes citrinus</u>	B
<u>Brachidontes exustus</u>	F, <u>exhustus</u> [sic] in ii, y
<u>Brachidontes recurvus</u>	B

<u>Brachidontes</u> sp.	jj
<u>Chama macerophylla</u>	KBM, B, +
<u>Chione cancellata</u>	x
<u>Chlamys imbricata</u>	+
<u>Codakia costata</u>	B, f
<u>Codakia orbicularis</u>	B, F, x
<u>Codakia orbiculata</u>	B, F, f, x, (as <u>Ctena</u> in ii), +
<u>Codakia pectinella</u>	x
<u>Coralliophaga coralliophaga</u>	B, f, +
<u>Corbula caribaea</u>	x
<u>Corbula contracta</u>	B, x
<u>Corbula cubaniana</u>	x
<u>Crassinella martinicensis</u>	JR, +
<u>Crassinella lunulata</u>	ii
<u>Crassostrea rhizophorae</u>	y
<u>Crassostrea</u> sp.	jj
<u>Ctena orbiculata</u>	ii
<u>Cumingia antillarum</u>	JR, B, x, +
<u>Cyathodonta semirugosa</u>	CB, +
<u>Diplodonta punctata</u>	B, F, x
<u>Diplodonta semiaspera</u>	JR, +
<u>Donax denticulatus</u>	CB, +
<u>Echinochama arcinella</u>	CB, +
<u>Erycina emmonsii</u>	B
<u>Erycina periscopiana</u>	B
<u>Gastrochaena hians</u>	B, f
<u>Gouldia cerina</u>	x
<u>Gregariella coralliophaga</u>	B
<u>Isognomon alatus</u>	CB, y, +
<u>Isognomon bicolor</u>	CB, B, F, +
<u>Isognomon radiatus</u>	CB, B, D, +
<u>Laevicardium laevigatum</u>	DLW, +
<u>Lima pellucida</u>	B
<u>Lima scabra</u>	DLW, +
<u>Lima scabra</u> form <u>tenera</u>	DLW, +

<u>Lioberus castaneus</u>	B
<u>Lithophaga antillarum</u>	KBM, f, +
<u>Lithophaga bisulcata</u>	B, F, f
<u>Lithophaga nigra</u>	B, F, f
<u>Lucina leucocyma</u>	x
<u>Lucina pensylvanica</u>	B
<u>Macoma constricta</u>	x
<u>Macoma tenta</u>	B
<u>Modiolus americanus</u>	KBM, B, +
<u>Musculus lateralis</u>	JR, f, +
<u>Ostrea equestris</u>	DLW, JR, +
<u>Phacoides muricatus</u>	x
<u>Phacoides pectinatus</u>	KBM, CB, B, F, ii, +
<u>Pinctada radiata</u>	KBM, B, F, +
<u>Pinna carneae</u>	JR, +
<u>Pseudochama arcinella</u>	B
<u>Pteria columbus</u>	DLW, KBM, JR, +
<u>Sphenia antillensis</u>	B, +
<u>Spondylus americanus</u>	SW
<u>Solemya sp. aff. occidentalis</u>	x
<u>Strigilla sp.</u>	B
<u>Tellina alternata</u>	ii
<u>Tellina exerythra</u>	x
<u>Tellina fausta</u>	DLW, JR, f, x, +
<u>Tellina listeri</u>	x
<u>Tellina nitens</u>	x
<u>Tellina promera</u>	ii
<u>Tellina versicolor</u>	x
<u>Tellina vespuciana</u>	x
<u>Teredo sp.</u>	y
<u>Trachycardium muricatum</u>	DLW, +
<u>Venericardia sp. aff. tridentata</u>	x

CEPHALOPODA

Octopus sp. E75, K

ECTOPROCTA (BRYOZOA)

<u>Arborella dichotoma</u>	JD, +
<u>Bugula</u> sp.	y
<u>Caberea carabaoda</u>	JD, +
<u>Caulibugula dendrograpta</u>	JD, +
<u>Celleporaria albirostris</u>	JD, +
<u>Chlidonia pyriformis</u>	JD, +
<u>Gemmillipora</u> sp.	y
<u>Gemelliporidra multilamellosa</u>	JD, aa, +
<u>Lichenopora buskiana</u>	JD, +
<u>Retoporelliria evelinae</u>	JD
<u>Steganoporella magnilabris</u>	JD, aa, +
<u>Stylopoma informata</u>	JD, +
<u>Trematooecia aviculifera</u>	JD, aa, +
<u>Trematooecia turrita</u>	JD, aa, +
<u>Tubucellaria cereoides</u>	JD, +

BRACHIOPODA

<u>Discinisca strigata</u>	B, + (specimen in collection from Pacific)
----------------------------	---

ECHINODERMATA

(by class)

ECHINOIDEA

<u>Arbacia punctulata</u>	DLM, +
<u>Brissopsis elongata</u>	DLM, +
<u>Brissus unicolor</u>	DLM, B, E76, k, +
<u>Diadema antillarum</u>	B, E74, E75, E76, k, o. v, x, :
<u>Echinometra lucunter</u>	B, E74, E75, E76, k, o, x

<u>Echinometra viridis</u>	B, E74, E75, E76, k, o
<u>Echinoneus cyclostomus</u>	B, E76, k, +
<u>Eucidaris tribuloides</u>	DLM, B, E76, k, o, y, +
<u>Halodeima floridana</u>	x
<u>Lytechinus variegatus</u>	DLM, B, E74, E75, E76, k, o, p, x, y, +
<u>Lytechinus williamsi</u>	DLM, +
<u>Meoma ventricosa</u>	DLM, +
<u>Paraster floridiensis</u>	DLM, E76, k, +
<u>Plagiobrissus grandis</u>	DLM, +
<u>Tripneustes ventricosus</u>	DLM, E76, k, v, x, +, <u>T. esculentus</u> in B

ASTEROIDEA

<u>Ophidiaster guildingii</u>	B
<u>Oreaster reticulatus</u>	DLM, B, +

OPHIUROIDEA

<u>Amphiodia repens</u>	B
<u>Amphiodia</u> sp. L	B
<u>Amphipholis</u> sp. A	B
<u>Amphipholis</u> sp.	B
<u>Amphipholis gracillima</u>	x
<u>Amphiura (Monamphiura)</u> sp.	B
<u>Amphiura (Nullamphiura)</u> sp.	B
<u>Axiognathus squamata</u>	B
<u>Ophiactis savignyi</u>	DLM, B, x, +
<u>Ophiocantha ophiactoides</u>	B
<u>Ophiocoma echinata</u>	DLM, B, +
<u>Ophiocoma pumila</u>	DLM, B, +
<u>Ophiocoma</u> sp.	y
<u>Ophiocoma wendti</u>	DLM, B, +
<u>Ophioderma appressum</u>	DLM, B, y, +

<u>Ophioderma brevicaudum</u>	DLM, B, +
<u>Ophioderma brevispinum</u>	DLM, B, +
<u>Ophioderma cinereum</u>	DLM, B, +
<u>Ophioderma rubicundum</u>	DLM, +
<u>Ophiolepis paucispina</u>	DLM, B, F, +
<u>Ophiomyxa flaccida</u>	DLM, +
<u>Ophionereis reticulata</u>	DLM, B, +
<u>Ophiothrix angulata</u>	DLM, B, +
<u>Ophiothrix cf. lineata</u>	DLM, +
<u>Ophiothrix suensonii</u>	DLM, +
<u>Ophiozona impressa</u>	DLM, +
<u>Ophiozonoida (?) sp.</u>	B (this may be new genus; EK, pers. comm.)

CRINOIDEA

<u>Comactinia echinoptera</u>	DLM
<u>Comactinia echinoptera:</u>	
cf. var. <u>valida</u>	cc
<u>Comactinia echinoptera</u> cf. var.	
<u>meridionalis</u>	cc
<u>Comactinia meridionalis</u>	DLM, +
<u>Nemaster rubiginosa</u>	cc
<u>Nemaster discoidea</u>	cc

HOLOTHUROIDEA

<u>Euapta lappa</u>	JC
<u>Holothuriid</u> sp.	JC

CHORDATA

ASCIDICEA

<u>Ascidia interrupta</u>	DS
---------------------------	----

<u>Clavelina</u> sp.	y
<u>Clavelina picta</u>	RHM
<u>Cystodytes dellechiajei</u>	RHM
<u>Didemnum</u> spp.	DS
<u>Didemnum</u> sp. B	RHM
<u>Didemnum</u> sp. C	RHM
<u>Didemnum</u> sp. D	RHM
? <u>Didemnum</u> sp. E	RHM
<u>Distaplia</u> (? <u>bermudensis</u>)	RHM
<u>Ecteinascidia</u> <u>conklini</u> var.	
<u>minuta</u>	DS
<u>Ecteinascidia</u> <u>turbinata</u>	RHM
<u>Eudistoma</u> sp. A	RHM
<u>Eudistoma</u> sp. B	RHM
<u>Pyrura</u> <u>vittata</u>	DS
<u>Polysyncraton</u> (? <u>amethysteum</u>)	RHM
<u>Rhopalaea</u> <u>abdominalis</u>	RHM
<u>Trididemnum</u> <u>cyanophorum</u>	
(= <u>T.</u> <u>solidum</u>)	RHM, RO, v, w
<u>Trididemnum</u> sp. A	RHM

NOTE: RHM identifications are provisional.

ACKNOWLEDGMENTS

We thank the following people for their assistance in preparing this species inventory: Victoria Batista, Charles Birkeland, Edwin Bourget, Leo Buss, Roy Caldwell, Hugh Dingle, Kristian Fauchald, Joan Ferraris, Robert H. Gore, Eugene Kaplan, Arilla Kourany, Haris Lessios, David Meyer, Kaniaulono Meyer, Randy Olson, James Porter, Klaus Ruetzler, Michael Robinson, Argelis Roman, Joseph Rosewater, Kenneth Sebens, Stephen Shuster, Richard Steger, and Janie Wulff.

This study was supported by the Smithsonian Institution Environmental Sciences Program and the Smithsonian Tropical Research Institute.

LITERATURE CITED

- Abele, Lawrence G. 1972. A review of the genus Ambidexter (Crustacea: Decapoda: Processidae) in Panama. Bull. Mar. Sci., 22(2): 365-380.
- Abrams, Peter A. 1976. Field guide to the hermit crabs of Galeta and vicinity. Smithsonian Tropical Research Institute. 4 p. (with subsequent revisions by PAA) unpubl.
- Batista, Victoria. 1980. Estudio de las comunidades que habitan las raices del mangle rojo Rhizophora mangle L. de Punta Galeta, Costa Atlantica de Panamá. Dissertation. Fundacion Universidad de Bogota Jorge Tadeo Lozano, Bogotá, Colombia.
- Bertsch, Hans. 1975a. Distributional and anatomical observations of Berthella tupala (Opisthobranchia: Notaspidea). The Nautilus, 89(4): 124-126.
- Bertsch, Hans. 1975b. Additional data for two dorid nudibranchs from the southern Caribbean seas. The Veliger, 17(4): 416-417.
- Birkeland, Charles. 1974. The effect of wave action on the population dynamics of Gorgonia ventalina Linnaeus. Studies in Tropical Oceanography, 12: 115-126.
- Birkeland, Charles, Amada A. Reimer and Joyce Redemske Young. 1976. Survey of marine communities in Panama and experiments with oil. EPA-600/3-76-028, National Technical Information Service, Springfield, VA 22161.
- Brattegård, Torleiv. 1974. Mysidacea from shallow water on the Caribbean coast of Panama. Sarsia, 57: 87-108.
- Caldwell, Gloria S. 1981. Attraction to tropical mixed-species heron flocks: proximate mechanism and consequences. Behavioral Ecology and Sociobiology, 8: 99-103.
- Child, C. Allan. 1979. Shallow-water Pycnogonida of the Isthmus of Panama and the coasts of Middle America. Smithsonian Contributions to Zoology, 293: 1-86.
- Connell, J. H. 1978. Diversity in tropical rain forests and coral reefs. Science, 199: 1302-1310.
- FAO. 1969. Symposium on investigations and resources of the Caribbean Sea and adjacent regions. FAO Fisheries Report No. 71.1.
- Fauchald, Kristian. 1977. Polychaetes from intertidal areas in Panama, with a review of previous shallow-water records. Smithsonian Contributions to Zoology, 221: 1-81.

- Gore, Robert H. 1976. Shallow water porcelain crabs from the Pacific coast of Panama and adjacent Caribbean waters (Crustacea: Anomura: Porcellanidae). Smithsonian Contributions to Zoology, 237: 1-30.
- Gore, Robert H. 1977. Neopisosoma angustifrons (Benedict, 1901): The complete larval development under laboratory conditions, with notes on larvae of the related genus Pachycheles (Decapoda Anomura, Porcellanidae). Crustaceana, 33(3): 284-300.
- Gore, Robert H. and Lawrence G. Abele. 1973. Three new species of Porcellanid crabs (Crustacea, Decapoda, Porcellanidae) from the Bay of Panama and adjacent Caribbean waters. Bull. Mar. Sci., 23(3): 559-573.
- Hendler, Gordon. 1977. The differential effects on seasonal stress and predation of the stability of reef-flat echinoid populations. Proceedings, Third International Coral Reef Symposium, Rosenstiel School of Marine and Atmospheric Science, Univ. of Miami, FL 33149, pp. 217-223.
- Henry, D. P., and P. A. McLaughlin. 1975. The barnacles of the Balanus amphitrite complex (Cirripedia, Thoracica). Zoologische Verhandelingen, 141: 1-254, + 22 plates.
- Jackson, J. B. 1973. The ecology of molluscs of Thalassia communities, Jamaica, West Indies. I. Distribution, environmental physiology, and ecology of common shallow-water species. Bulletin of Marine Science, 23: 313-350.
- Koehl, M. A. R. 1977. Water flow and the morphology of zoanthid colonies. Proceedings, Third International Coral Reef Symposium, Rosenstiel School of Marine and Atmospheric Science, Univ. of Miami, FL 33149, pp. 437-444.
- Lasker, Howard R. 1979. Light dependent activity patterns among reef corals: Montastrea cavernosa. Biol. Bull., 156: 196-211.
- Lawrence, John M. 1976. Covering response in sea urchins. Nature, 262: 490-491.
- Lehman, John T. and James W. Porter. 1973. Chemical activation of feeding in the Caribbean reef-building coral Montastrea cavernosa. Biol. Bull., 145: 140-149.
- Macintyre, I. G. and P. W. Glynn. 1976. Evolution of modern Caribbean fringing reef, Galeta Point, Panama. Am. Assoc. Petrol. Geol. Bull., 60(7): 1054-1072.
- Meyer, D. L. 1973. Feeding behavior and ecology of shallow-water unstalked crinoids (Echinodermata) in the Caribbean Sea. Mar. Biol., 22: 105-129.

- Meyer, Kaniaulono B. 1977. Dorid nudibranchs of the Caribbean coast of the Panama Canal Zone. Bull. Mar. Sci., 27(2): 299-307.
- Neumann, C. J., G. W. Cry, E. L. Caso, B. R. Jarvinen. 1978. Tropical cyclones of the North Atlantic Ocean, 1871-1977. U.S. Department of Commerce, NOAA, National Weather Service, Environmental Data Service.
- Olson, Randy. 1979. An ecological investigation of Trididemnum cyanophorum (Lafargue and Duclaux, 1979): a compound tunicate with symbiotic algae. STRI Short-term Fellowship Unpublished Report.
- Porter, James W. 1972. Ecology and species diversity of coral reefs on opposite sides of the isthmus of Panama. Bull. Biol. Soc. Wash., 2: 89-116.
- Powell, N. A. 1971. The marine bryozoa near the Panama Canal. Bull. Mar. Sci., 21(3): 766-778.
- Reimer, Amada A. 1975. Effects of crude oil on the feeding behavior of the zoanthid Palythoa variabilis. Environ. Physiol. Biochem., 5: 258-266.
- Ruetzler, K., and W. Sterrer. 1970. Damage observed in tropical communities along the Atlantic seaboard of Panama. BioScience, 20: 222-224.
- Sebens, Kenneth P. 1976. The ecology of Caribbean sea anemones in Panama: utilization of space on a coral reef. In G. O. Mackie (ed.), Coelenterate Ecology and Behavior, Plenum Pub. Corp., New York, pp. 67-77.
- Sebens, Kenneth P. 1977. Autotrophic and heterotrophic nutrition of coral reef zoanthids. Proceedings, Third International Coral Reef Symposium, Rosenstiel School of Marine and Atmospheric Science, Univ. of Miami, FL 33149, pp. 397-404.
- Sebens, Kenneth P. 1982. Intertidal distribution of zoanthids on the Caribbean coast of Panama: Effects of predation and desiccation. Bulletin of Marine Science, 321: 316-335.
- Sebens, K. P. and K. DeRiemer. 1977. Diel cycles of expansion and contraction in coral reef Anthozoans. Mar. Biol., 43: 247-256.
- Smith, Ralph I. and James T. Carlton (eds.). 1975. Light's Manual: Intertidal Invertebrates of the Central California Coast, 3d ed. Univ. of California Press. 716 p.
- Smithsonian Institution Environmental Sciences Program. 1974. Environmental Monitoring and Baseline Data 1973. Tropical Studies. Roberta W. Rubinoff (ed.). Smithsonian Institution, Washington, DC.

- Smithsonian Institution Environmental Sciences Program. 1975.
Environmental Monitoring and Baseline Data 1974. Tropical Studies.
Donald M. Windsor (ed.). Smithsonian Institution, Washington, DC.
- Smithsonian Institution Environmental Sciences Program. 1976.
Environmental Monitoring and Baseline Data 1975. Tropical Studies.
Donald M. Windsor (ed.). Smithsonian Institution, Washington, DC.
- Smithsonian Institution Environmental Sciences Program. 1977.
Environmental Monitoring and Baseline Data 1976. Tropical Studies.
Donald M. Windsor (ed.). Smithsonian Institution, Washington, DC.
- Smith, Wayne L. 1973. Record of a fish associated with a Caribbean sea anemone. Copeia, 1973(3): 597-598.
- Southward, A. J. 1975. Intertidal and shallow water Cirripedia of the Caribbean. Studies on the fauna of Curacao and other Caribbean islands, 46: 1-53.
- Southward, A. J., and W. A. Newman. 1977. Aspects of the ecology and biogeography of the intertidal and shallow-water balanomorph cirripedia of the Caribbean and adjacent sea-areas. FAO Fisheries Reports, 200: 407-425.
- Spivey, H. R. 1976. The Cirripeds of the Panama Canal. Corrosion and Marine-fouling, 1: 43-50.
- UNEP (United Nations Environment Programme). 1981. Intergovernmental meeting on the action plan for the Caribbean Environment Programme. Report of the meeting. UNEP/CEPAL/IG.27/3.
- Vasquez-Montoya, Rafael. 1979. Peuplements des herbiers de Thalassia testudinum et d'Halodule sp. de la Côte Caraïbe de Panama. Ph.D. These. A. L'Université D'Aix-Marseille II. 40 p. + Appendices.
- Yee, Gloria B. de and Carmen Chang. 1978. Mortalidad de los Erizos en la Costa Atlantica de Panama. ConCiencia, 5(1): 7-8.